

Publicaties

- Blokhorst MGBG, Slot L & Winter F (1994). *Het Brein de Baas; het zelf behandelen van geheugenproblemen*. Soest: Ruitenbergboek.
- Slot L & Blokhorst MGBG (1995). *Slapen leer je zo*. Uitgeverij: Van Holkema & Warendorf.
- Blokhorst MGBG, Lousberg R, Vingerhoets A, Winter F, Zilvold G. (2002a). Daily Hassles and Stress-vulnerability in patients with Whiplash Associated Disorder. *International Journal of Rehabilitation Research*, 25, 173–179.
- Blokhorst MGBG, Swinkels M, Lof O, Lousberg R, Zilvold G. (2002b). The influence of state related factors on focused attention following Whiplash Associated Disorder. *Journal of Clinical and Experimental Neuropsychology*, 24, 471–478.
- van Toor T, Neijenhuis K, Snik A & Blokhorst MGBG. (2003). Evaluation of Auditory processing disorders after whiplash. In: *Auditory Processing Disorders. Development and evaluation of a test battery*. (K. Neijenhuis, Thesis). Nijmegen: Radboud University, 109–129.
- Wassink B, Somer R & Blokhorst MGBG (2004). Strategietraining in een revalidatiecentrum. *Neuropraxis*, 8(1), 28–31.
- Blokhorst MGBG, Meeldijk S, van Luijtelaar G, van Toor T, Lousberg R. (2005a). Noise intolerance and state-dependent factors in patients with whiplash associated disorder. *Journal of Whiplash and Related Disorders*, 4,(1).
- Blokhorst MGBG, Schrijver P, Meeldijk S, Hermans R, Lousberg R. & Zilvold G. (2005b). Cortisol Responses to experimental stress in patients with Whiplash Associated Disorder. *'Stress and Health' - new research*. New York: Nova Sciences.
- Blokhorst MGBG. (2005c). *State-dependent factors and attention in Whiplash Associated Disorder*. Thesis. Enschede, UT/RRD.
- Blokhorst MGBG & Schreurs K (2006). Whiplash en stress: het Biopsychosociaal Whiplashmodel. *Stimulus*, 4, 233–256.
- Wouters EJM, Reimus JLM, van Nunen AMA, Blokhorst MGBG & Vingerhoets AJM (2008). Suffering Quantified? Feasibility and Psychometric characteristics of 2 revised versions of the pictorial representation of Illness and Self measure (PRISM). *Behavioral Medicine*, 34, 65–76.
- Blokhorst MGBG (2011). Wanneer is cognitieve gedragstherapie bij rugklachten geïndiceerd? *Physios*, 4, 24–30.